



The Spire Sentinel



**The Newsletter & Magazine of The
Chesterfield Branch of The Western Front
Association**



ISSUE 82 - December 2022

Our aims are 'Remembrance and Sharing the History of the
Great War'.



Western Front Association Chesterfield Branch – Meetings 2022

Meetings start at 7.30pm and take place at the Labour Club, Unity House, Saltergate, Chesterfield S40 1NF

January	4th	.Branch AGM and Members Evening - 3 short presentations by Jon-Paul Harding, Andy Rawson and Grant Cullen
February	1st	`Steaming to The Front` - Britain`s Railways in The Great War by Grant Cullen
March	1st	`They Think It`s all Over` By Andy Rawson . Plenty has been said about the breaking of the Hindenburg Line. This presentation looks at the pursuit of the Germans which occurred during the final weeks of the war.
April	5th	Soldiers and Their Horses - Horses and Their Soldiers by Dr Jane Flynn - a sympathetic consideration of the soldier - horse relationship 1914-18
May	3rd	`Finding Deborah` by Mike Tipping. How the team that discovered tank Deborah D-51 went on to find me, and my journey to Deborah
June	7th	The Cost of the War' By Roy Larkin. Hansard tells us that the Great War of 1914-18 increased the National Debt to £7,435,000,000 or £377,144,063,927 at today's value which took 100 years to pay off.
July	5th	The Italian Front 1915-1918 by John Chester. Covers the fighting in Italy from beginning to end. Includes the contribution of the British and their part in ending the war.
August	2nd	Peter Hart returns to Chesterfield - last time was just before the first lockdown in March 2020. The title of Peter`s talk is Rupert Brooke and the `Glitterati` at Gallipoli`
September	6th	The Inventions Department by Richard Godber. A little known part of the Ministry of Munitions. Based upon Richard`s dissertation for his Wolverhampton MA, previously a very under researched area about which little was known.
October	4th	`The Fighting Fifth`and the attack at Bellewarde Ridge 16th June 1915 by John Beech. John has a strong personal connection with the Northumberland Fusiliers in this action
November	1st	`Shell Shock and the History of Psychiatry` by Jill Brunt. Based upon sessions on this subject presented to students at Northern College, Barnsley
December	6th	From Sheffield to Serre - The City Battalion at Home by Andy Rawson - complimenting the guided trips to the sites of Redmires Camp earlier this year

Issue 82 - list of contents

- 2 Branch Meetings Calendar
- 3 Contents Page + Monday Night Webinars + Events for 2023
- 4 Secretary`s Scribbles
- 5 Branded Goods
- 6 WFA 2023 Calendars
- 7 - 8 Service of Remembrance at Cenotaph November 11th 2022
- 9 - 16 November Meeting
- 16 - 27 Terror Weapons: The British Experience of Gas and Its Treatment in the First World War

WFA Webinars have now recommenced, under is a list of December 2022 Webinars, please register if you are interested in watching,

12 DEC 2022 ONLINE: From Plumstead to Palestine – Some Cockney War Stories

<https://www.westernfrontassociation.com/events/online-from-plumstead-to-palestine-some-cockney-war-stories/>

19 DEC 2022 ONLINE: The Russian Civil War and the Allied Intervention Force

<https://www.westernfrontassociation.com/events/online-the-russian-civil-war-and-the-allied-intervention-force/>

2023 Programme of Talks – January – May

January 3rd AGM + *‘British League of Help’* by Dudley Giles. Nearly 90 towns, cities, and organisations in the UK, Australia, Canada and Mauritius signed up in the period 1920-1922 to 'adopt' a village, town or city in the Devastated Zone of France. Some of these adoptions lasted only a few years, some (like Sheffield's adoption of Bapaume, Serre and Puisieux) survived until after WW2

February 7th *‘The First AIR War’* by Grant Cullen. Based on a collection of rare photographs acquired over 20 years ago at a yard sale in Hazelwood, Missouri, US, this will look at the various protagonists in WW1 - people and planes - finishing with a detailed look at the rotary aero engine and the air mounted Lewis gun

March 7th - *‘Voie Sacree’* by Roy Larkin. The story of the road that connects Bar-le-Duc to Verdun It was given its name because of the vital role it played during the Battle of Verdun in World War I.

April 4th *‘For Home and Honour’* by Yvonne Ridgeway and James Kay. A bit of a history of our local community in North Sheffield during WW1, from their own research, looking at recruitment, the 1st Sheffield blitz, the tribunals for those wishing to avoid military service and some of the local soldiers' stories.

May 2nd *The First World War contribution of Dulmial Village , present day Pakistan* by Dr Irfan Malik. His Gt. Grandfathers experiences in WW1, and the wider role of muslim soldiers during that conflict

Any opinions expressed in this Newsletter / Magazine are not necessarily those of the Western Front Association, Chesterfield Branch, in particular, or the Western Front Association in general



Secretary's Scribbles

Dear Members and Friends,

Welcome to the last edition of our Branch Newsletter for 2022.

As reported elsewhere Jane Lovatt, Jon-Paul Harding, Yvonne Ridgeway and James Kay went to the ceremony at the Cenotaph in London on November 11th. All the costs being met by Branch funds plus a generous grant given to Branches by WFA Central for this purpose.

Of course these guys were not the only representatives of our Branch being in attendance. Tony Bolton was there as National Chair and Rob Nash who assisted John Chester as Parade Marshall. The ceremony was live streamed for the first time.

As previously advised due to unforeseen circumstances our scheduled speaker for December is unavailable - that talk will now take place in January 2023.

I am very grateful for Andy Rawson taking over the December slot - his talk is quite apt given he led two `expeditions` to the Redmires site a couple of months back, anyway here is the details....

From Sheffield to Serre - The City Battalion at Home

It covers the story of the recruitment and early training of the Sheffield City Battalion through the stories which appeared in the newspapers, using the many photographs they used at the time. Andy will be looking at what happened through the eyes of the people, rather than the historians. It means he focuses on the battalion's experience and reactions to it between September 1914 and May 1915, when they headed off to Cannock Chase, en route to Egypt and the Somme. It is also a good insight into the early days of a Pals Battalion. Andy has a bit of a personal connection - his grandparents lived in the two farms overlooking the camp.

All coming along next Tuesday meeting will be invited to stay for some `nibbles` and a free drink in the Labour Club bar after the conclusion of Andy`s presentation. It`s our way of saying `thank you` to those who have supported us during 2022 which has seen us bounce back to pre covid attendance levels.

The January presentation will be `*British League of Help*` by Dudley Giles postponed from December.

I`m still working to produce a full calendar of events for next years programme but the first five months of 2023 are fixed up - details elsewhere - I`m waiting confirmation from other speakers to whom invitations have been sent out .

Best wishes, Grant Cullen Branch Secretary 07824628638



BRANDED GOODS AVAILABILITY

New items are always being considered, so please check the Branded goods part of the shop for all items available.

Prices are inclusive of postage within UK (Branded Items Nos 1-11)

www.westernfrontassociation.com/shop/wfa-branded-items/?p=2

or call Head Office (Sarah Gunn or Maya Shapland) on 020 7118 1914

And the (Branded Clothing, Nos 12- 18) note new prices (under) effective from 1st July.

Order direct from supplier (West Coast Workwear) www.westernfrontassociation.com/shop/branded-clothing/ or ring (0800 169 2228 or 01704 873301)



- | | | | |
|----|-------------------|----------|--|
| 1 | Fridge Magnet | (£5) | 59mm dia, front metal plate, high strength neodymium magnetic backplate, and plastic mylar front cover |
| 2 | Anniv' Coaster | (£8.50) | 4" in diameter hand crafted slate. Individually polished, screen printed by hand and backed by a baize |
| 3 | Mousemats | (£6) | 196 x 235mm fabric surface and are of high quality. They have a rubberised base layer |
| 4 | Bookmarks | (£2) | (dims 55 x 175mm) rich UV High Gloss Coating provides protection against stains and damage |
| 5 | Baseball Caps | (£8) | Lightweight 5 Panel cotton cap, adjustable with velcro rip-strip, one size fits all |
| 6 | Ties | (£11) | Length 142cm, width 9cm (at widest part), 100% Polyester |
| 7 | Lapel Badges | (£2.50) | 25mm Dia. Die struck + imitation hard enamel, Silver Nickel Plating, Butterfly clutch pin |
| 8 | Mug | (£10) | 11oz ceramic mug (95mm high x 85mm diameter) features the bold official WFA logo design (two sides) |
| 9 | Messenger Bag | (£27) | 37 x 29 x 11cm, 100% Cotton. Full cotton lining. Zippered organiser section, Capacity:13 litres |
| 10 | Despatch Bag | (£30) | 40 x 30 x 12 cm, (10) Washed Canvas, dual rear pouch pockets. Multiple zippered pockets. Capacity: 14 litres |
| 11 | Shoulder Bag | (£25) | 40 x 28 x 18 cm, (10) (11) Polyester. Internal valuables pocket. Zippered front pocket. Capacity: 14 litres |
| 12 | Oxford Shirt | (£27) | Kustom Kit Short Sleeve Corporate Oxford Shirt. Easy iron button down collar, 85% cotton, 15% polyester |
| 13 | Breathable Jacket | (£71) | Russell Hydro Plus 2000 Jacket. Nylon taslon with PU Coating |
| 14 | Rugby Shirt | (£25) | Front Row Classic Rugby Shirt, 100% Cotton |
| 15 | Fleece | (£24) | Regatta Thor 111 Fleece Jacket, 100% polyester anti pill |
| 16 | T-shirt | (£17) | Russell Classic Cotton T-Shirt. 100% ringspun cotton |
| 17 | Sweat Shirt | (£22.50) | Gents Russell Jerzees Raglan / Ladies Fruit Of The Loom Raglan |
| 18 | Polo Shirt | (£20.50) | Russell Cotton Pique Polo Shirt. 100% cotton |

WFA 2023 CALENDARS

The Western Front Association's 2023 calendar is now available. Once again it features images of the battlefield taken by a team of volunteer photographers. The scenes depict points of interest in France and Belgium (and, incidentally, Italy) some of which are well known but others 'off the beaten track'. As well as providing superb images of a dozen views of the First World War battlefields, the calendar provides detailed commentary to each image helping to set the scene in context. This is a high-quality product which, every year, receives superb feedback. The sales of the calendar also assist the WFA to continue its work. The WFA's 2023 calendar is available via [the WFA e-shop](#) or by phone on 0207 118 1914.

A few of the images are shown below:



WFA Chesterfield - Deputation to the Service of Remembrance at The Cenotaph, London, November 11th 2022. Report by Jane Lovatt

With the impetus of a grant from the WFA, Jon-Paul, Yvonne, James and I decided to make the journey to London to attend the 11 November Cenotaph ceremony. After a ridiculously early start from Chesterfield....we arrived ridiculously early! Skirting around Buckingham Palace we drove down an almost deserted Mall. Deputy Parade Marshall of the event, and Chesterfield Branch regular Rob Nash had kindly arranged a pass which enabled us to park yards from the Cenotaph on King Charles Street. After clearing security, officials took pity on the unlikely group of visitors and ushered us into the Foreign Office to use the 'facilities'. An auspicious start to the day.

Fortunately London was enjoying an almost spring-like day and, with the excitement of children on a school visit we, decided to do some sightseeing. Turning left out of King Charles Street, we approached the Cenotaph and Downing Street. A group of veterans being ushered into the Prime Minister's residence caught my eye. At this point the Downing Street Police security detail having spotted our poppies, asked us for details of the Cenotaph ceremony. To our astonishment they informed us that, although they were aware that the ceremony was taking place, they were ignorant of the details and asked us if we knew the timings.

We took advantage of a relatively quiet Whitehall to take photos and enjoy the monuments and memorials scattered along the route to Trafalgar Square. Here we grabbed a very welcome coffee before heading up the Strand and taking a circuitous route back to King Charles Street.



Jon-Paul was delighted to be nominated to lay a wreath on behalf of the Branch and was ushered away by Phylomena Badsey. Shortly before 11am the excited atmosphere changed to a more serious note as we formed up and processed on to a cordoned off Whitehall. A sense of stillness and a dignified silence descended on this busy part of the Capital. Wreath layers lined up flanking the front of the Cenotaph with a party of dignitaries to the side. Our Branch Chairman and WFA National Chairman Tony Bolton heading up the party. The ceremony proceeded with practised precision under the watchful eyes of John Chester and Rob Nash. The lone bugler of the Grenadier Guards and the bandsmen of the London Scottish lent a poignant air to the occasion. A sizeable crowd of onlookers had gathered to join with the ceremony and pay their respects.

With military precision wreaths were laid and speeches made under the ever watchful eye of Earl Haig who is memorialised yards from the Cenotaph.

Following the ceremony we made our way to The Royal Military Chapel-The Guards Chapel- for the service of remembrance. It was an honour to be in the presence of the colours and standards that adorn the Chapel walls. The silver candles which survived the Second World War bombing being perpetually lit. The talented choir and sermon by Reverend Morgan were interspersed with readings by WFA members. The service concluded with the Last Post performed by the Grenadier Guards, one minute's silence and Reveille.



The Chesterfield WFA contingent - left to right - Yvonne Ridgeway, James Kay, Jane Lovatt, Jon-Paul Harding.

Afterwards we meandered back to the car and took the decision to head home before the Friday evening rush.

We all enjoyed, and felt privileged to have had the opportunity to remember and celebrate our ancestors at the National service.

November Meeting

Branch Chair (and National Chair !) Tony Bolton welcomed everyone, remarking on the excellent attendance - the best since before covid restrictions. It was great to see branch stalwarts Arthur Lacey and Alan Atkinson back among us for the first time since early 2020.

Tony then introduced our speaker for the evening Jill Brunt.



Jill started her working life as a Speech and Language Therapist, practicing in the NHS and treating both adults and children. A change in career and a Masters Degree led her in to Adult and University Education. She was Assistant Principal at Northern College, Senior Research Fellow and Director of Teaching in the Faculty of Medicine, University of Sheffield and CEO of an Awarding Body. Jill held two Visiting Professorships at Sheffield Hallam University and London South Bank University.

She is now a Chesterfield Borough Councillor with a Cabinet Portfolio for Health and Well-being.

Jill`s talk was entitled ` **Shell Shock in WW1** ` and would explore the position of psychiatry during the war, presenting symptoms and treatment for shell shock and the context in the UK for returning soldiers.

Jill started by explaining how she became interested in WW1, by visiting the battlefields with husband Steve, reading books and watching films, particularly the film `Regeneration` about failure and inability to communicate. It is very important to understand the historical context in dealing with mental health, what systems were in place to deal with `shell shock` when confronted with it, what happened when soldiers returned home, a little bit about women and finally a bit about what Jill described as the `other` war dead.

Psychiatry during the war could not be relied upon as a source of comfort, shell shock exposed an area of medicine about which most asylum doctors knew very little. From the military standpoint, a deserter was `considered insane and destined for the madhouse or irresponsible and should be shot` a comment written by a Colonel Myers, more about him later.

So therefore things were not in a readable state to understand shell shock and how it could be treated. We were in a period which came to be known as the `Great Lock Up`. Before 1914 insanity was all about finding people who were insane and locking them up in asylums and just leaving them there. The whole issue of combat brought a very different challenge for mental health professionals used to dealing with insanity in civilian life.

The psychosomatic disorders brought new military-medico terms for them, things like `soldiers heart` and of course we already used terms like `bad with the nerves`. But there was also a feeling that by 1900 psychiatry had reached a dead end only concentrating on asylums and people detained there because of this great `Lock Up`...get `lunatics` off the streets just lock them away. There was so many of them that any sense of treatment, recovery or discharge was really just a lost hope.

From the literature of the time it can be seen that psychiatrists of the day were looked upon as `second rate` within the medical professionals..a bit like the old saying...`those who can do...those who can't ..teach...`. In other words if you were not going to be a good doctor then you might as well be a psychiatrist!

Those who are intellectually inadequate instinctively seek out an asylum where they could become superintendents there. That`s what people thought about the profession.

By 1914 the Medico-Psychological Association of Great Britain and Ireland stated that psychiatry as a profession needed reform but within a few weeks of that report being published, war broke out and nothing further was done. The absence of early provision of early diagnosis and treatment of mental illness was seriously under-developed and there was a linguistic failure as well, a failure to describe...what words could be used. There was a lack of facilities to research these conditions and a very poor provision of assistant medical officers in asylums, people were admitted to asylums and there was inadequate treatment in these places. Also living conditions, food etc., were really poor. There was a notion that our country had lagged behind other civilised nations into the treatment of mental illness. If you think of Freud later on in 1914 much of the rest of Europe was moving on further than we were in terms of treatment., especially in Russia who had witnessed the experiences of soldiers from their war with Japan in 1905 and had begun to change some of their services accordingly.

When the hospitals of Europe started to take their first cases of shell shock the British Medical Journal of that time suggested that treating drunkenness and syphilis was considered more important. The stress of war started to make some people think in a very different light and to seek much more rational and humane treatments for cases of mental disorder. The wider community was obsessed with eugenics - once a lunatic always a lunatic...or his dad / mum was a lunatic or `it runs in the family` The public took little interest in psychological disorders and there was little attempt to understand the condition of these patients. In some of the literature there was harsh condemnation of those who tried to understand mental health - issues - you were a soldier first and a medic second - patients were not first.

The average training of a medical student in mental health involved only one visit to an asylum where they saw the very advanced stages of mental illness and this was probably a very traumatic experience for some of these medical students to have to go into an asylum when you read of the conditions in these places at that time - they were horrendous.

The attention given to mental diseases before qualification was much less than given to other countries, again, we were lagging behind. Owing to the lack of mental

health clinics students had no opportunity to observe borderline or underdeveloped cases so looking at the notion of the `journey` of mental illness was just not available, they were just thrown in at the deep end or only able to look at what was going on in asylums.

There was very few teaching facilities in asylums or what was then termed the Lunacy Service and the training that was given to doctors was quite hopeless. In Germany patients were free to come and go, there was early detection, less stigma and psychiatric clinics attached to every university. There was a much larger ratio of doctors and nurses to patients. In Britain, asylums were not in a good shape at all - one medical officer to between 400 to 600 patients. Today the ratio of GPs to patients is about 2000 as Arthur Lacey commented.

Very much a result of the Mental Health Act of 1890-91 which delayed the diagnosis and as we know early diagnosis and intervention in any health issue is key to recovery. The Lunacy Law prevented the establishment of psychiatric clinics as they had in Germany and people had to be certified as a lunatic and sent to an asylum. There was no change in recovery rates from the Crimean War to World War One, in other words you went in, weren't going to recover and never come out. At the same time there was improvement in diphtheria, heart disease and tuberculosis but the absence of research into mental illness meant there was little improvement there. Again, one of the views in the literature was that the asylum system or Great Lock Up was to separate lunatics from the gene pool of the nation.



Many pictures of the time show soldier suffering Shell Shock` to have the `thousand yard stare` of showing little behind the eyes

There was a notion that those with shell shock were emotionally incontinent...they laughed or cried at the wrong time

There were some unexplained physical manifestations, people walking strangely, losing the feeling in their legs or facial tics.

Now the term that shell shock would be called severe neurosis as a result of being under fire as, particularly on the Western Front, soldiers were subjected to a seemingly incessant barrage of being under shellfire. Tiredness, irritability, giddiness, lack of concentration, headaches, tremors, mutism and special twitches were all seen to be symptoms of shell shock. There was many ideas about what actually caused

those symptoms and an early thought was that those bursting shells created a vacuum and when the air rushed into the vacuum it upset the cerebro-spinal fluid, this upsetting the working of the brain. They were trying to find real reasons why this should be happening. There was some references from the American Civil War - `soldiers` heart` etc. fatigue, lack of energy etc. as they tried to identify the causes and from there define some kind of treatment. In Vietnam it was combat fatigue and in the Gulf War PTSD. There have been a lot of improvements and training in understanding the generic `shell shock`

The language that was used in WW1... `he`s lost his mind...he`s lost his reason...he`s bad with his nerves` ...sometimes the patient himself thought he was mad, being left alone and told to cheer up, didn`t help . There was an exaggeration of emotion, the exposure to cold, wet, and hunger conditions only served to make things worse.

Another symptom was to appear to his comrades to be less fearful of danger, taking unnecessary risks as seen in the film `Regeneration` and there are many stories like this from the battlefields. There were attempts to suppress emotions but the physical symptoms, the breathing and digestive disorders. Suppression of emotions can be exhausting - something we all know from our own experiences - trying to hide your feelings which actually makes it much worse in the long run.

Often soldiers would arrive in the clearing hospitals with what was known as `delusional insanity` and recovered quickly, a fact which made diagnosis really difficult. There is a common thread running through all the shell shock literature as to how you determine whether somebody was a coward or a weakling or were genuinely mentally disturbed. Often patients would be in hospitals were they were not encouraged to talk which led them to believe they were uncared for and beyond help. Chloroform was used to treat mutism as was electric shock applied to vocal chords - absolutely barbaric. Some clinicians were applying these techniques while others therapeutic therapies and towards end of WW1 communities were being developed and this was continued into the 1920s, one of them being the York Retreat developed from the English Quaker community both as a reaction against the harsh, inhumane treatment common to other asylums of that era, and as a model of Quaker therapeutic beliefs. A common belief at the time was that the mad were wild beasts. The recommended medical practices included debilitating purges, painful blistering, and long-term immobilisation by manacles, and sudden immersion in cold baths - all administered in regimes of fear, terror and brutality. The Retreat continued to operate as an independent hospital into the modern era. However, on 31 December 2018, it withdrew from inpatient care with its remaining services for eating disorders and personality disorders now being run by the Clinic. The Retreat continues to run outpatient community psychological assessment, diagnostic and therapy services at the Tuke Centre, including an Autism and ADHD service. An M.O. Col. Myers based in St. Omer was filtering cases who would otherwise have been sent back to the front to the asylum. He believed that shell shock was a distinctive category of nervous disorder that was not lunacy. He developed a category of `NYD` - Not Yet Diagnosed` which allowed patients more time to recover. He was fighting against the military - medical community who considered shell shock to be an exit ticket for dirty sneaks and weaklings. In the UK upon arrival at the asylum they would be stripped of their uniform straight away as they were considered to be a disgrace

to it. If someone was kind to them they would be in civvy suit so has not to have to undergo the shame of stripping down.

The treatment of shell shock mirrored all the divisions of a pre-war society - officers treated very differently from Other Ranks, for example at Craiglockhart inmates were allowed to move around quite freely vastly different from the asylums where in 1919 former soldiers who had been incarcerated in asylums were complaining of starvation. The Poor Law Infirmary or the Workhouse was another option but irrespective they were not well looked after.

It was an achievement in itself to survive in the asylum, some inmates actually starving to death and there was always the risk of falling into the hands of the `bodysnatchers` of which there were many at that time and the lunatic asylums were a fertile source of bodies. In subsequent years trying to find records of inmates was difficult, with records often being incomplete or in some cases destroyed. Those who returned to work and were still troubled with their nerves did not find favour with the Ministry of Pensions. At the end of the war in 1918 shell shock `exploded` onto the streets of Britain with sufferers walking the streets and this led to general questioning as to what was going on and the country became very close to having another `Great Lock Up`. Cases of domestic violence, assault, delusions shot up and there is also the factor that some people went into the war with a mental health problem and came out of it with a similar condition.

There was questions as to whether inmates of asylums suffering from shell shock should form the Community of Remembrance at The Cenotaph they being viewed as an embarrassment with no recognition that they too, had served their country.


The Ministry of Pensions issued guidance as to how service patients should be buried if they escaped incarceration in the asylum as a corpse - should they be buried in uniform should anyone in uniform attend the burial - they were considered a disgrace.

The psychiatric aftermath of the Great War was becoming increasingly difficult and public opinion was gathering momentum as neurasthenia and hysteria was presenting itself after discharge. Treatment was only given to patients on the condition that they entered asylum thereby declaring themselves as insane and giving themselves over to custody. For a soldier coming back from the war with serious mental health issues yet having a wife and children to support, the only way they could get money was to enter an asylum.

The National Federation of Disabled and Discharged Soldiers approached the Prime Minister asking that a nationwide network of specialist institutions be set up to treat soldiers with shell shock where they could get the appropriate treatment not available in the lunatic asylums.

By 1920, the military establishment tried to distance itself from the psychiatric issues arising from the war, to instil into the public a more traditional moral condition, to put `Tommy` in his place and represent shell shock to the nation and there followed from that a whole series of events and conferences saying that shell shock didn't really exist and that it was just a lot of people `swinging the lead` or being cowardly. This affected people's pensions or their ability to get work.

Shell shock is a confusing term which did not help anyone differentiate from those who were having a mental breakdown, men who were simply exhausted and those who were, in fact, `swinging the lead`. There was overuse of the term without proper diagnosis. Jill recommended the book Broken Men, for those interested in further information.



Broken Men: Shell Shock, Treatment And Recovery In Britain 1914-30: Shell Shock, Treatment and Recovery in Britain 1914-1930

As Jill moved on, she that that concluded her overview of how people with shell shock were treated.

In terms of literature it is difficult to find out much about women in the First World War - except for the VADs which of course stood for Very Adorable Darlings - but Jill did pick out two. First, Dame Maude Macarthy



McCarthy sailed in the first ship to leave England with members of the British Expeditionary Force (BEF), arriving in France on 12 August 1914. In 1915 she was installed at Abbeville as matron-in-chief of the BEF in France and Flanders, taking charge of the whole area from the Channel to the Mediterranean, wherever British and allied nurses worked; she was directly responsible to General Headquarters. In August 1914, the numbers in her charge were 516; by the time of the Armistice they were over 6,000. She was responsible for the nursing of hundreds of thousands of casualties from 1914 to 1918.

She was one of the first to recognise shell shock – but her name does not appear in any contemporary shell shock literature. She lobbied the various nursing institutions to give the VADs recognition of the training they received and again – what we would call `recognition of accredited learning`

Dr Isobel Stoneham served in military hospitals in France , Malta and Egypt a rarity as women doctors - and there was less than 1% of all doctors being women - were told

to `stay at home` and take up the slack caused by male doctors leaving to go to the front. Also, when they did appear in other areas of the world they were not allowed to wear a uniform although that was not the case in French Red Cross hospitals.

However, by the summer of 1916 more doctors were needed and they were allowed to work alongside RAMC staff in Malta...they could work in a French hospital and Malta...but not in the English system. Male doctors who volunteered received temporary commissions entitling them to wear uniform but not female doctors. The Medical Women`s Federation campaigned and the army reluctantly permitted them to wear uniforms but commissions for women required legislation.

The `Other` War Dead. In late 1914 there was a recognition that there was not enough hospital beds to cope with the growing number of casualties. Recovering soldiers needed both beds and open space recreational areas and the lunatic asylums as they were called then could offer those facilities. Every county in England had a least one asylum and a number of workhouses. There was 102 asylums for 108000 men, women and children living there permanently, some having lived there for a very long time. Living conditions for people in these places with mental illness, Downes Syndrome - those with that condition were put in asylums - epilepsy, T.B., alcoholism...again reflecting the great lock up...let`s just lock up people away from the rest of society. The Asylum Act 1915-1919 saw the emptying of all the asylums which were swiftly renamed war hospitals indeed within 5 weeks all the selected asylums had been emptied of their inmates, only the dying and those termed `quiet, useful`...those who could worked for example in the gardens, remained.

The whole gamut of emotions was exhibited by the patients on leaving ranging from acute distress and misery, indifference as to what was happening to them to maniacal fury and indignation. This reported by the Medical Officer at Norfolk County Asylum. Basically they had been living there for years that was their home and most had no understanding as to what was happening to them. Asylum patients died at a greater rate after leaving the hospital than previously....from TB, `Flu etc.

By the 1920s over 440000 soldiers, from all over the world had been treated in these former asylums with 38000 being shell shock cases. A subsequent enquiry reported that the transferred insane should be viewed as quasi-casualties of the war and that had never happened. The hidden cost was that over 12000 of the most vulnerable people in Britain were forcibly displaced.

There are a number of contested areas - firstly the number of shell shock cases, it was said that there was 80000 cases of shell shock but this has been contested by many people who consider that 250000 would be a more accurate figure...who counts - what about those who recovered quickly. The Military Medical Statistics ended the day that war was over so those who returned after that date were described as `injured` but not recorded as having shell shock. The Military Medical Officers at the end of the war were more or less working for the Ministry of Pensions - not for the patients. The country could ill afford pensions awarded to shell shock victims. Are you a medic first and soldier second...or a soldier first and a medic second?. That of course also applied to chaplains, Jill saying that husband Steve had recently acquired a very good book called `The Flag` ...again what are you - chaplain first soldier second or soldier first - chaplain second...?

For Jill the overwhelming issue was the state of the mental health system at the time and the state of psychiatry and how could it have been managed differently and as she said, look at the progress since then, talking therapy...the Freudian ways of the 1930s...the lessening of stigma today surrounding mental health.

That ended Jill`s formal presentation but it didn`t end the meeting as we had a prolonged discussion, with many attendees contributing to what turned out to be a very informative.

Once all the Q & A was over, Branch Chair Tony Bolton thanked Jill for an informative talk which had become such an interesting event proposing a vote of thanks to which the attendees warmly responded.

Terror Weapons: The British Experience of Gas and Its Treatment in the First World War



Edgar Jones

King's College London, UK

Abstract

Chemical weapons accounted for only 1 per cent of the 750,000 British troops killed in the First World War and yet caused disproportionate casualties (estimated at 180,100). The considerable investment in the development of new toxins and methods of delivery was designed to maintain the elements of surprise and uncertainty as these accentuated their psychological effect. Soldiers were continually challenged on the battlefield by combinations of different types of agent designed to undermine their confidence in respirators, disorientate them, and erode their morale. At first army doctors practised defensive medicine, invaliding their patients for protracted periods to the UK or base hospitals. By 1917, progressive study of the physical and psychological effects of different types of toxin allowed physicians to design new management strategies. Borrowing ideas from shell shock, specialist units were set up closer to the front line and medical officers taught to identify crucial points in the course of illness to accelerate recovery times and forestall the accretion of psychosomatic symptoms.

Chemical weapons and observed, 'gas shock was as frequent as shell shock'.

The psychological impact of these toxins was confirmed by Lieutenant Colonel C. Gordon Douglas, a physiologist and specialist gas officer, who concluded that 'the particular value of the poison [mustard gas] is to be found in its remarkable casualty producing power as opposed to its killing power'. The capacity of gas to inspire fear was apparent from its first large-scale use on the Western Front. At Langemarck, on 22 April 1915, the release of 150 tons of chlorine from 6,000 cylinders caused widespread panic. The chaotic retreat of two French divisions, 87th Territorial and 45th Algerian, opened a 4 mile gap in the front line. As these troops had no protective equipment or any training in gas warfare, it was scarcely surprising that they fled when confronted with a suffocating, greenish-yellow cloud. In September 1915, when the British released chlorine in retaliation, similar effects were observed of German troops at Loos:

A German officer in this sector remarked that as soon as the gas entered his trench, he lost all control of his men, a panic ensued and he was unable to keep them in the front line. He said that, without the gas, we should have had no earthly chance of taking the trenches. Uncontrolled anxiety during a gas attack could cause men to tear off their protective masks, or act 'as though they had temporarily lost their reason'. Later in the war Charles Wilson, a regimental medical officer with the Royal Fusiliers, argued that mustard gas had 'partly usurped the role of high explosive in bringing to a head a natural unfitness for war, or less commonly in undermining fitness sapped by exceptional stress in the field'.

Terms such as 'gas hysteria' and 'gas neurosis' were coined to describe enduring somatic symptoms once physical lesions had healed. Hulbert thought 'gas neurosis' akin to shell shock of a non-concussive type because the severity of symptoms bore no relation to 'the amount of gas inhaled' but arose 'in proportion to the individual's mental and emotional make-up and instability'.

While the initial psychological impact of gas was explicable in terms of surprise and lack of preparedness, its enduring effectiveness as a terror weapon requires explanation. Almost 60 per cent of deaths in the First World War were a result of artillery and trench mortars; by comparison, gas killed few troops. Furthermore, most soldiers exposed to chemical weapons survived and made a good recovery. In a post-war study Brigadier Augustin Prentiss of the American Chemical Warfare Service estimated that only 4.3 per cent of gas casualties died, compared with 24 per cent of other types of battlefield injury. From mid-1916, respirators offered troops reliable protection against chlorine and phosgene. Yet gas remained among the most feared weapons of the war and continued to exercise a powerful hold over the popular imagination such that anti-war campaigners focused on its use to mobilize support for their cause.

Edward Spiers and L.F. Haber have documented the scientific resources deployed to devise and identify new chemical weapons, together with innovative ways of manufacturing and delivering these toxins. In addition, Donald Richter has explored the heroic efforts devoted to finding protective measures and training soldiers in their use. Helen McCartney discovered that witnessing the effects of chlorine was a defining moment for the Territorials of the Liverpool Scottish as it brought home the horrors and inhumanity of war. In his study of morale in the German and British Armies, Alexander Watson argued that gas created uncertainty: unlike shrapnel, it killed from the inside, eroding a soldier's sense of control, while raising the terrifying fear of being suffocated. As regards the treatment of gas casualties, Mark Harrison has explored the evolution of their management within the provision of medical care in France. Spiers has also looked at the narratives of gas in the post-war period and how these fed into popular culture and the political debate about the future use of chemical weapons. However, with the exception of Tim Cook's study of Canadian gas services, less research has focused on the psychological impact of gassing. This paper explores how different groups of soldiers responded, how symptoms developed over time, and the investigations conducted by medical officers to improve diagnosis and

treatment. It analyses the effectiveness of the managerial strategy of the British Expeditionary Force (BEF) to reduce the amount of time that gassed soldiers spent in hospital and increase the proportion that returned to active duty.

I. Physical Effects of Gas

Although chlorine was readily detected by its pungent odour and yellow-green clouds, phosgene was more difficult to identify, being colourless and having the smell of freshly cut hay. In December 1915, for example, the Germans introduced phosgene, which was six times more potent than chlorine and could be inhaled in fatal doses without the coughing and discomfort associated with chlorine. Furthermore, the symptoms of phosgene could be delayed for several hours, making immediate diagnosis problematic. Indeed, it was estimated to have caused 85 per cent of all deaths from chemical weapons during the First World War. To distinguish between the unpleasant (tear gas and the chlorarsines) and the lethal (chlorine, phosgene, and mustard gas) took nerve and training. Indeed, chlorarsines caused short-term but intense respiratory distress designed to disable temporarily but also to terrify. Colonel A. Bertram Soltau, consultant physician in France for gas cases, emphasized the importance of chemical weapons in the genesis of 'nervous disorders': there 'is nothing', he argued, 'probably more liable to cause panic than the idea of being choked ... the dread of being slowly strangled'.

The surprise element, so important for gas, was maintained throughout the war by the continual development of new toxins and delivery methods. When outlining British policy in June 1916, Field Marshal John French had argued that it was 'essential that the nature of the gases discharged from cylinders and in projectiles should be varied from time to time in order that the enemy's protective measures may be rendered as difficult as possible'. Planners deliberately exploited ignorance and fear of chemical weapons. For example, an infantry assault on the British First Army at Vermelles on 27 April 1916 was preceded by lachrymatory gas shells and clouds of harmless smoke. Before the second attack, an hour later, lethal chlorine was released, designed to surprise soldiers who had removed their respirators. Used against the British from July 1916 onwards, the gas shell enabled a range of toxins to be delivered rapidly without warning. In addition the British produced the Livens projector, which, according to captured German troops, was the most demoralizing weapon the Allies possessed, making life 'utterly unbearable'. First used at the battle of Arras in April 1917, the Livens projector propelled a drum containing 30 lb of chemical (usually phosgene) over a range of 1,700 yards. In batteries of 25 they delivered a massive quantity of agent with accuracy and little warning. In response the Germans devised a 'Gaswerfer', which gave a high concentration of lethal gas by firing hundreds of large phosgene projectiles over a limited front.

In 1976 Ludwig Haber estimated that 6,060 soldiers of the BEF had died as a result of gas (about 1 per cent) and a further 180,100 (3.3 per cent) had been injured. Official statistics compiled by Mitchell and Smith recorded 185,706 gas casualties admitted to hospitals in France, with 5,899 deaths. According to this data, admissions rose steadily throughout the war; those for 1918 (113,764) were almost double those for 1915-17 (71,942). However, this temporal rise was primarily a function of increased use, the quantity of chemical consumed having risen from 3,870 tons in 1915 to 65,160 tons in 1918. Greater numbers of patients did not represent a widespread failure of treatment or protective measures. In fact, deaths fell as a proportion of admissions from 4.5(3,226) for 1915-17 to 2.3 (2,673) in the last year of the war. Reduced mortality was a consequence of better quality respirators and improved clinical techniques, notably oxygen therapy.

At first British medical authorities had struggled to contain the problem. Not only did numbers killed by cloud gas rise, death rates at casualty clearing stations (CCSs) rose progressively from 3.6 per cent in December 1915 to 19.6 per cent by August 1916

This data, collected by Major T.R. Elliott and Captain C.G Douglas in an attempt to understand the impact of chemical weapons, established that gas could be a deadly and potent weapon. The rising mortality that they documented was a consequence of increased use of phosgene, difficulties in developing an effective respirator, and inadequate training of medical staff in diagnosis and treatment. Indeed, the statistics related to a time when Royal Army Medical Corps (RAMC) physicians believed that most gassed patients would benefit from bleeding (venesection) to reduce strain on the heart and drain fluid from the lungs. Not until 1917 was it established that the procedure was actually harmful in cases characterized by a rapid and weak pulse.

If worn correctly, the small box respirator, introduced by the British late in 1916 (though not fully issued until January 1917), provided reliable protection against chlorine and phosgene. Largely for this reason, in July 1917 the Germans introduced mustard gas, a vesicant that burned the skin. Known as 'yellow cross' from the shell markings, the toxin immediately produced a flood of admissions: 160,970 soldiers presented at CCSs between 21 July and 23 November 1917, of whom 1,859 died. It was estimated that 75 per cent had been exposed to mustard gas. Deaths initially occurred because the toxin had delayed effects. The smell of the gas was 'not very noticeable and ... the immediate effect on troops exposed to it', Brigadier K. Wigram reported, was 'only a slight irritation of the nose and throat', so that most troops had been slow to realize that they had been poisoned. Mustard gas achieved its greatest effect in the months immediately after its introduction because 'the novelty of the condition, the multiplicity of the symptoms, and John Singer Sargent's large canvas 'Gassed' depicted a corps dressing station at Le-Bacde-Sud on the Doullens road. Soldiers arrived in groups suffering from the effects of gas shells, while others lay on the ground. Exhibited at the Royal Academy in spring 1919, 'Gassed' became an iconic image of the suffering caused by the war. In Germany, Otto Dix used the image of the soldier wearing a respirator to symbolize the inhumanity of combat and to show soldiers as instruments of an industrial conflict.

The entire absence of knowledge as to possible after-effects naturally led to the condition being overestimated'. Fear inspired by mustard gas was heightened by its capacity to cause loss of sight. Although most blinded servicemen recovered, the acute photophobia, conjunctivitis, and oedema of the eyelids forced soldiers to close their eyes, 'so much so in fact that when some of the milder cases were evacuated each man had to be led like a blind man by an orderly to the ambulance car'. Such images struck a cultural chord and remain among the iconic symbols of the First World War.

Data collected by Douglas towards the end of the war, and subsequently published in the official history, demonstrated that mortality rates were determined by the nature of the toxin, delivery method, and effectiveness of protective measures. Although cylinder gas was subject to the vagaries of the prevailing wind, its use corresponded with a time of inadequate respirators and little anti-gas training. As a result, it generated high mortality rates. By contrast, mustard gas, deployed after the issue of the box respirator when seasoned troops had habituated to the threat of chemical weapons, led to much lower death rates, even though its delivery was far more precise. Consequently, death rates were as low as 2.5 per 100 casualties for mustard gas and as high as 24.0 for troops exposed to chlorine and phosgene in the period before the introduction of the small box respirator. Interestingly, phosgene delivered by gas shell caused more casualties, but a significantly lower death rate, than cloud gas. The reduced mortality was a consequence of greatly improved protective measures, confidence in their use, and the growing provision of specialist treatment. The high death rate for projector attacks related to their capacity to deliver high concentrations of deadly agent with minimal warning. The physical effects of poison gas, though sometimes terrifying to observe, were in the majority of cases not fatal and most soldiers made a good recovery. If, however, a man received a lethal dose of chlorine or phosgene, death commonly came within two days. By comparison, mustard gas, first used against British troops on the night of 12/13 July 1917 at Ypres, was designed to

disable rather than kill. It was estimated that of the 180,100 British chemical casualties, 120,000 had been subjected to mustard gas.

Provided a soldier wore a gas mask correctly, his respiratory system was protected, but woollen uniforms offered no defence against blistering of the skin. Research showed that a thick suit coated with oxidized oils and resins neutralized the vesicant effects of mustard gas, but such a garment was 'hot and uncomfortable to wear, and for a fully equipped soldier marching and fighting in such clothing would be impossible'. Although gas capes were issued to British troops during the Second World War, no corresponding protection was forthcoming in 1917-18. American medics and gunners were, however, issued with an 'anti-gas suit' made of cotton sheeting impregnated with linseed or vegetable drying oil.

II. Psychological Effects

Even before the mass use of chemical weapons on the battlefield, the 'subjective effect' of toxins on an individual's mind had been recorded. Early in 1915 British scientists tested the possible use of ethyl iodoacetate (a lachrymator given the code name 'South Kensington' after the experimental work conducted at Imperial College of Science and Technology). A number of army officers from Chatham garrison were invited to attend field trials: 'One of them, who was stationed at least 50 yards up wind from the point of burst, immediately left the trench showing every sign of great mental disturbance and stating that he felt very ill.' It was established that he could not have inhaled any of the vapour and yet had been deeply affected by the experience.

Douglas observed that, although not primarily designed to inspire terror, the 'violent irritant or choking sensations' of chlorine and phosgene had the capacity to undermine the resolve of all but the most resolute soldier. Specialist medical officers increasingly recognized the importance of gas as a psychological weapon. Captain H.W. Barber, who treated mustard gas cases at No. 25 General Hospital, argued that the 'sudden shock' of being gassed often caused as many symptoms as 'any toxic property of the gas itself'. Writing in spring 1917, Lieutenant Colonel S.L. Cummins, adviser in pathology to the British armies in France, concluded that any division subjected to a series of gas attacks in close succession was likely to exhibit a significant drop in morale, while Charles Wilson, a regimental medical officer, believed that 'the majority of men who left the front line in 1917 "gassed" were frankly frightened'.

The capacity of poison gas to inspire strong emotion led to a range of unwanted outcomes: panic even when protected by a respirator, the misinterpretation of harmless sounds and smells and taking evasive action, soldiers reporting sick when actually well, and doctors referring mild or transient cases of gassing for lengthy treatment in base hospitals. Panic is defined as precipitate and unreasoning behaviour not likely to serve the interests of the subject; it often involves actual or attempted physical flight. Captain A.J. Waugh, medical officer to the 1st North Staffords, reported such a case when his battalion was exposed to cloud gas in May 1916: 'a few men lost their heads, took off their [anti-gas] helmets and ran back, being severely gassed in consequence'. Examples of troops misinterpreting harmless visual and olfactory stimuli were common and revealed the power of gas to disrupt military routine and discipline. Lieutenant G.L. Grant, medical officer of the London Scottish, treated large numbers of officers and men in September 1915 who believed they had been gassed. None had any objective signs of toxic exposure and all responded miraculously to a placebo. In February 1918 a soldier in a working party of 1/22nd London Regiment felt swelling and soreness of his throat and reported that he had been gassed. Although no shelling had taken place and no one had observed any signs of gas, fear swept through the unit and within a few hours of the 105 men had been evacuated to an advanced dressing station as casualties, where some 'acted as though they had temporarily lost their reason'. No organic cause could be discovered and the fact that no officer reported any ill effects suggested that this was an example of contagious anxiety. Similarly, a group of US machine-gunners became

convinced that their food had been contaminated by toxins from a shell that had exploded nearby. Presenting to a nearby aid post, they complained of stomach pains, and some had even vomited. Doctors could find no evidence of exposure to gas and they were treated with bicarbonate of soda.

Diphenyl chlorarsine ('blue cross'), a nasal irritant and vomiting agent first used against the British in July 1917, caused short-term incapacity: sometimes 'the pain and discomfort is sufficient to cause a man to lose his mental control for a short time'. Although the toxic effects were temporary, servicemen often continued to experience symptoms after the poison had worn off. Five soldiers were examined by Captain C.D. Christie three days after they had been exposed to chlorarsine gas. They complained of 'extreme weakness and inability to use all of their extremities'. Christie observed that 'it is very hard to reconcile the bizarre nature and distribution of the neurological findings on any anatomical or physiological basis', though he believed the symptoms to be 'genuine', which suggested an unconscious mechanism rather than malingering.

III. Adaptation and Protective Measures

Although gas masks saved lives by offering reliable protection against the inhalation of toxins, in themselves they were also a source of anxiety for both wearers and onlookers. Captain G. Donaldson of the 2/7th Royal Warwickshire Regiment, writing home in July 1916, observed: 'We had our gas helmets on. It was like an appalling night-mare as you look like some horrible kind of demon or goblin in these masks.' When 'smoke and tube helmets' were first introduced in autumn 1915, some units discarded them, having misinterpreted the smell of neutralizing chemical with which they were impregnated for that of gas; the practice helmets that they had worn earlier had little or no odour. Habituation and the adoption of coping strategies were hampered by continual refinements in chemical weapon technology. Each toxin had specific properties, demanding different protective measures and different forms of treatment. Knowledge, even among the medical corps, remained inconsistent. As late as April 1918 Douglas observed, 'I really believe that nearly all medical officers are terrified of the mere mention of gas poisoning,' and a month earlier had acknowledged that 'the majority of medical officers' in France could not accurately diagnose which gas a patient had been exposed to on the basis of its physical effects. Captain W.J. Forshaw, based at No. 2 Australian General Hospital, wrote of mustard gas in 1917: 'many regimental medical officers have no knowledge of the after effects and receive no information and scanty supplies of material for treatment'. Doctors untrained in the effects of gas practised defensive medicine and referred patients to base hospitals whether or not this caution was needed. Knowledge can, but does not necessarily, serve as an effective defence against irrational fear. However, mastering protective measures and diagnosing the differences between different types of gas took considerable training. The use of phosgene against British troops on 19 December 1915 prompted the setting up of anti-gas schools to prepare soldiers for the hazards of chemical warfare, and from March 1916 instruction in anti-gas measures to recently arrived drafts was given at Étaples, Rouen, and Havre. In September an anti-gas school also opened at Calais and in the following year others at Boulogne and Abbeville. Soldiers were exposed to an hour of cloud gas to give them confidence in their mask, and then exposed to 30 seconds of tear gas to give them a fright. Contemporary reports conflicted on the effectiveness of this training. Some servicemen reported a harsh realism. Private Frank Bass of the 1st Battalion, Cambridgeshire Regiment, wrote of his time at Étaples in September 1916: 'Lecture on gas. Officer lecturing had been two years here and through two gas attacks. Callousness of lecturers shocks us.' Many gas instruction officers felt devalued and, despite the dangers of chemical weapons, found soldiers unmotivated by the subject. A.E. Hodgkin, chemical adviser to the Fifth Army, recalled that many slept through his lectures unless compelled by the cold to stay awake: 'never, never will the mystery of gas warfare penetrate the brain of the regular soldier'. More effective, perhaps, was the experience of seeing fellow soldiers poisoned by gas. Ernst Jünger recalled that sight of comrades 'groaning

and retching while their eyes watered' taught him 'never to go anywhere without my gas mask, having previously, incredibly foolishly, often left it behind in my dugout, and used its case - like a botanist - as a container for sandwiches'. However, contemporary accounts suggested that well-disciplined and experienced troops sustained lower rates of gas casualties than battalions newly arrived at the front.

Douglas observed in March 1918:

We have admittedly to deal with a large psychic element in very mild cases of gas poisoning, and this feature is naturally more prominent in the case of troops who have only a limited experience of gas shelling. Even with such experienced troops of our own, we have to contend with this difficulty - hence instructions to detain doubtful cases of gas poisoning in medical units within the army area until diagnosis is certain.

The official history reported that when exposed to chlorine at Bellewaarde Ridge on 24 May 1915, soldiers of the 5th Border Regiment, attached to 10th Brigade, showed no outward concern. Many were miners accustomed to the dangers of gas, and their example did much to 'fortify the confidence of other troops'.⁸⁵ Such variations in responses to chemical attacks confirmed that some units habituated to the threats; the question remains whether those battalions that accommodated more readily had been better prepared in terms of training and selection for the demands of trench warfare. No studies were undertaken to test the value of anti-gas training and the impact of actual exposure on subsequent performance.

IV. Management Strategies

The management of gas casualties (that is, systems introduced to direct the flow of patients from the battlefield through various treatment processes to duty or discharge from the forces) evolved during the conflict as a small number of specialist physicians acquired technical understanding. C.G. Douglas was a key player in the BEF's strategy to combat the effects of gas. A physiologist and fellow of St John's College, Oxford, he had volunteered for military service in 1914 and was sent to France, where, in May 1915 at the suggestion of J.S. Haldane, he was transferred from general medical duties to investigate the immediate effects of chlorine on front-line soldiers. Douglas had conducted research into respiratory physiology under Haldane's guidance and was the obvious person to study the pathology of chemical weapons. He visited the sites of gas attacks and studied casualties in detail, building up a comprehensive understanding of their diagnosis and treatment. Indeed, medical officers in command of CCSs were ordered to inform Douglas whenever severely gassed cases were admitted to their units so that he could make an immediate examination. Awarded a Military Cross in 1916 and four mentions in dispatches, Douglas rose to the rank of lieutenant colonel in 1918 and was responsible for drafting the British army's official policy on the effects of gas and their treatment. On being demobilized from the forces, he returned to his Oxford laboratory and a distinguished career of academic research in which he collaborated with J.G. Priestley to write a textbook, *Human Physiology*. His precision, attention to detail, and personal courage were in no small measure responsible for the accretion of reliable data on which to base practice and policy.

After the first major use of gas in 1915, and without an obvious treatment apart from bed rest, the British medical service in France adopted a defensive policy: the transfer of gas casualties to the UK as quickly as possible. This management approach became deeply embedded in RAMC culture, despite mounting evidence to suggest that it was often inappropriate and inefficient. The general medical strategy of 1915-16 was to build up a network of CCSs with a range of expertise to take the pressure off base hospitals during offensives. In terms of gas casualties, the CCSs were designed to filter out mild cases to prevent them overcrowding treatment facilities in the rear, but because doctors in these units had only rudimentary knowledge of chemical weapons, they referred the vast majority of such patients to base hospitals.

By dramatically increasing the number of casualties, the use of mustard gas in July 1917 forced the British to re-evaluate the way that servicemen were treated. Early in 1918 Douglas discovered that 58 per cent of gas casualties admitted to all base hospitals in France between 1 July and 31 October 1917 had been evacuated to the UK. Of those that had remained in France, only 23.3 per cent had returned to active duty by the beginning of November, while 17.5 per cent were still being treated in base hospitals or convalescent depots. The management of shell shock provided a model that could be adapted for gassed patients. In December 1916 forward psychiatric units had been set up in CCSs to offer rapid treatment and prevent evacuation to the rear or the UK.⁹⁵ Faced with a flood of mustard gas cases, the BEF adopted a similar strategy. Plans to open a specialist gas hospital for each of the armies in France were abandoned, 'as the object is to bring cases under treatment as early as possible'. As with shell shock, specialist officers were deployed to undertake diagnosis, and regimental medical officers were instructed to label gassed soldiers as 'Not Yet Diagnosed Gas' to pre-empt premature evacuation to base hospitals.

In practice the new policy proved difficult to implement. As late as March 1918, Douglas bemoaned 'the tendency to send gas casualties, even when they are very slight, over to England, and this of course implies very heavy wastage'. In September 1918 Colonel Elliott was horrified to discover that mustard gas cases were still being transferred to the UK, and only 11 per cent had returned to duty with the BEF after five months of treatment. Not only did this policy extend the period that soldiers remained patients, clinical studies conducted towards the end of 1916 showed that excessive periods of treatment could arrest a natural recovery process as new clusters of functional symptoms developed. By autumn 1918, surveys had shown that 70 per cent of mustard gas casualties treated in CCSs and other lines of communication units could be returned to duty within eight weeks of exposure.

V. Treatment Tactics

The first chlorine attacks put doctors in a difficult position. Few had any knowledge of how to treat the toxin, and medicine could offer little to counteract severe pulmonary damage. As a result, military physicians took an exceptionally precautionary approach to any case of poisoning. Great emphasis was placed on rest, and Sir Arthur Sloggett, the director general of army medical services, ordered that for a minimum of two days 'no casualty should be allowed to leave his bed or stretcher for any purpose whatever'. Severe cases were retained at CCSs for at least four days and transferred to base hospitals lying down. A report compiled in autumn 1916 showed that cases of moderate exposure were commonly held in medical units for two to three months 'and very possibly longer'. While this cautious approach had merit in spring 1915, when knowledge was incomplete, its value soon expired. For the majority of casualties, who proved to be mild cases, this management instilled or reinforced the idea that they were suffering from a potent and irreversible affliction.

Rather than promote an atmosphere of recovery and health, extended stay in hospital created fertile conditions for the elaboration of symptoms and chronic invalidity. Diagnostic difficulties created by the use of novel toxins were compounded because the symptoms of acute anxiety often mimicked the physical effects of mild exposure to gas: palpitations, chest pain, shortness of breath, fatigue, and weakness could all be produced by fear and worry, a similarity sometimes exploited by front-line troops seeking a medical exit from the battlefield. Having been invalided to a place of safety, the anxious soldier often made a recovery only to relapse when faced with the prospect of discharge from hospital and return to the trenches.

A survey conducted at No. 25 General Hospital at Hardelot, near Boulogne (which had a specialist gas ward), of 496 chlorine and phosgene casualties admitted between 8 July and 12 September 1916 produced disappointing findings. Only 118 patients (23.7 per cent) were discharged directly to base details and from there to active duty, while a further 132 (26.5 per cent) were referred to No. 1 and No. 5 Convalescent

Depots at Boulogne. By the end of September only 42 (8.5 per cent) of these convalescent servicemen had returned to duty. In total 179 (36 per cent) gassed soldiers had been evacuated to the UK for further treatment. Although the study had shown that two-thirds of cases could 'be satisfactorily treated in France', it also demonstrated that doctors had yet to identify clinical regimes that efficiently returned servicemen to active duty.

By May 1916 the threat to manpower presented by chemical weapons had become so serious that Sloggett set up a 'committee on treatment of gas cases'.¹¹⁰ With Cummins as its secretary, it comprised a group of specialist physicians: Elliott, Douglas, Sir Wilmot Herringham, and Sir Almroth Wright. The last, a bacteriologist, had worked with the British army to develop a vaccine against typhoid, and was in France to conduct research into wound infections. In an era before antibiotics, gassed soldiers were also vulnerable to infection. At the very outset the committee made a crucial error: 'in view of the fact that the cases arriving at base hospitals during the recent attacks have been, for the most part, slight or convalescent, it was decided that no evidence need be taken on the lines of communication'. By concentrating the study at base hospitals and excluding CCSs and other intermediate treatment centres, the committee failed to observe the development of symptoms and identify opportunities for early intervention. Not until 1917 was the oversight corrected.

Research into patterns of illness following exposure to gas revealed a complex picture. A study of convalescent depots in France by Captain Riddell in summer 1916 identified the features of disordered action of the heart (DAH) in gassed soldiers 'under protracted' treatment.¹¹³ DAH was characterized by shortness of breath, palpitations, chest pain, and fatigue after any form of exercise. No organic cause could be found and yet the disorder could prevent a soldier from returning to active duty. Furthermore, a study conducted in May and June 1916 of five convalescent depots receiving gassed casualties revealed extended treatment times: of 676 admissions, 480 (71 per cent) were retained for an average of nine weeks before being found 'fit for duty'. Further research at Mount Vernon Military Heart Hospital in Hampstead and at No. 25 General Hospital in France revealed that 'cardiac disability' was the 'chief weakness which invades all these [chlorine and phosgene] casualties'. At first, because of the mortality associated with severe exposure, doctors were misled by these symptoms. In June 1916 T.R. Elliott examined a selection of gassed DAH patients and concluded that they had either been prematurely encouraged to undertake programmes of graduated exercise or they had inhaled greater concentrations of gas than realized. Accordingly, Elliott recommended an extended period of recuperation and advised that any patient who exhibited an irregular heartbeat in the third week of admission should be transferred to the UK 'for rest'.

Yet the solution proposed by Elliott was soon shown to exacerbate the ongoing invalidity. During 1917 further study of gassed servicemen found that the acute effects of gas could pass only to be replaced by a range of psychosomatic symptoms (irregular heartbeat, chest pain, and shortness of breath), aggravated when asked to perform any form of physical effort. Specialist gas doctors largely agreed that this was not a toxic effect. Lieutenant Colonel W.E. Hume, who had studied mustard gas patients at No. 25 General Hospital and No. 1 Convalescent Depot at Boulogne, argued, 'the fact that there is such a discrepancy between the fast [heart] rate in all conditions of the body awake and the slow rate asleep in the majority of DAH patients seems to be proof that the tachycardia is of psychological origin'. In a report for the Medical Research Committee, J.S. Haldane, J.C. Meakins, and J.G. Priestley observed that 'it is difficult in many cases to distinguish the chronic gas cases from those suffering from irritable heart, shell-shock, or neurasthenia'. Similarly, Soltau, who examined the files of 150 gas pensioners, concluded that 30 per cent reported a range of symptoms that could equally well meet the criteria for neurasthenia or shell shock, while a further 25 per cent could be reclassified

as DAH, formerly known as soldier's heart.

Working at No. 15 Canadian General Hospital in Taplow, Buckinghamshire, Lieutenant Colonel John C. Meakins and Captain T.W. Walker studied chronic cases of gassing who had been invalided to the UK. The key symptom, they identified, was shallow breathing which appeared to prevent the patient from increasing the volume of their respirations beyond a limited extent. This, in turn, restricted any physical activity that they could undertake. Dyspnoea, or shortness of breath, was often accompanied by rapid heartbeat, dizziness, and fatigue. These symptoms were not necessarily correlated with severity of exposure to toxin and, in their view, reflected 'a distinct neurotic element'. The 'mental effect of gas poisoning', Meakins and Walker believed, was heightened by 'the delayed action of certain gases and frightful consequences of high concentrations ... deeply impressed upon the minds of the soldiers by the observation of their more unfortunate comrades'. A follow-up study conducted three months after discharge showed that only 16 (9.8 per cent) of 163 mustard gas cases returned to duty with the BEF.

Concerned by lengthy stays in medical units and the need to return as many soldiers as possible to fighting units, a group of RAMC doctors (C.G. Douglas, T.R. Elliott, and A.B. Soltau) decided to monitor admissions to find ways of making treatments more effective. They studied the progress of gassed patients as they passed from field ambulance to CCS, base hospital, and convalescent depot to find out more about lengths of admission and outcomes. Key points in treatment were identified when so-called 'neurasthenic' symptoms might develop. 'The neurasthenic element', Douglas argued, was the important feature in all gas cases, and it was the recognition of the true part played by this that contributed to the results. Firm control of patients from the start and the careful restriction of the period of detention in hospital to the minimum, prevented cases from falling into a morbid condition and developing those functional symptoms which so often delay convalescence and are exaggerated by prolonged hospitalization.

The doctors discovered considerable variation in the efficacy of treatment: only 19.4 per cent of gassed patients treated at No. 25 General Hospital, with its specialist gas ward, were referred to the UK, whereas at Étapes, which lacked such expertise, 62.1 per cent were invalided across the Channel.¹²⁵ From his study of No. 7 Stationary Hospital, Boulogne, Elliott concluded that 'no casualties need be invalided for a longer period than three months and that the majority are soon fit for duty'.

Following these investigations, strategies were developed to maintain the momentum of recovery and to distract patients from their symptoms. For example, men wearing dark glasses to combat the effects of photophobia were ordered to remove them as soon as their pupils had returned to a normal size and colour. When the acute effects of gassing had passed, great emphasis was placed on fresh air, exercise, and soldierly activities to maintain the progression to active duty. Programmes of graduated exercise were devised, based on measurements of pulse and respiration. The guidelines were: four days after the patient is allowed out of bed, he walks half a mile and, if this is not found excessive, he walks one mile on the next day; and if this again is not too much, it is repeated, and on the following day he walks three miles at the rate of three miles an hour. It continued to be believed that if a soldier were started on a programme of 'muscular exercise' too early or too forcibly, then tachycardia and dyspnoea were aggravated and the condition of DAH was 'liable to be established and to persist for a very long time'. Nevertheless, by mid-1918 Douglas estimated that the average treatment time for 80 per cent of gas admissions had been cut from around three months to eight weeks. Casualties, of whatever nature, invalided to hospitals in Britain were less likely to return to front-line battalions than those treated in France. This was not simply because they included the most severe cases but also because soldiers were understandably reluctant to surrender a place of safety for the dangers and privations of trench warfare. As a result, considerable resources were directed towards treating the wounded and sick in France, and specialist units for shell shock, functional heart disorders, and gassed servicemen

were set up. In October 1917, for example, a field ambulance serving the 47th Division was designated a corps gas centre not only to provide expert diagnosis and treatment but also to reduce the flow of casualties across the Channel. When, in August 1917, Douglas investigated the returns for all British base hospitals in France, he found that 47 per cent of gassed patients had been invalided to the UK for further treatment. Shows that the proportion evacuated home fell progressively from 22 per cent in summer 1918 to 5 per cent by the autumn. In addition, the specialist gas unit opened at No. 7 Stationary Hospital, Boulogne, under Major Wilson and Captain McIntosh steadily reduced the length of time spent in hospital. 'The experience of this hospital showed very clearly the value of special knowledge of gas poisoning on the part of the medical officer in order that he can be confident and firm in dealing with gassed soldiers.'

Tables also shows that convalescent depots opened on the French coast were increasingly used to prepare soldiers for return to active duty. With shorter times in hospital, they were designed, as Major General Sir Wilmot Herringham recalled, 'to give the men a cheerful and enjoyable time, while strengthening their bodies by regular and at the same time interesting exercise'. However, evidence from the convalescent depots suggests that it was far from easy to encourage soldiers to leave the relative comforts of these camps for the privations of the front line. In summer 1917 Douglas conducted a study of convalescent cases in France and found that after eight weeks only 64.7 per cent had discharged to full duty. Continued training of medical officers in diagnosis, together with stricter limits on periods of convalescence, progressively reduced the time that gassed soldiers remained as invalids. By September 1918, Colonel T.R. Elliott argued, this system of specialist hospital treatment combined with formal convalescence enabled the military to return 70 per cent of men exposed to mustard gas to infantry bases within eight weeks of exposure.

A study of ex-servicemen who had been awarded a war pension for gassing showed that, though they had recovered from the physical effects, many continued to suffer from medically unexplained symptoms and psychological effects. Many of these pensioners were reclassified as suffering from DAH because cardiac symptoms (palpitations, chest pain, and shortness of breath) were prominent. Much effort was devoted to their treatment, largely involving programmes of graduated exercise. Although oxygen therapy saved the lives of many with severe exposure to gas, 141 experiments showed that it had no lasting or therapeutic effect on chronic cases reclassified as DAH.¹⁴² Because of popular sympathy and the fact that many were unable to work consistently, many received a war pension.

VI. Conclusions

Basil Liddell Hart, who had himself been gassed at the Somme in July 1916, wrote, 'the chlorine gas originally used was undeniably cruel, but no worse than the frequent effect of shell or bayonet, and when it was succeeded by improved forms of gas both experience and statistics proved it the least inhumane of modern weapons'.¹⁴³ Although his classification of chemical weapons did not gain general currency in the post-war period, his observation about their capacity to kill or wound requires discussion. While poets such as Wilfred Owen emphasized the trauma of soldiers dying from gas, their suffering was not significantly different from a terminal stomach wound or shrapnel damage to the head and face. This raises the question whether gas had a particular capacity to inspire terror, or whether the initial novelty and the continual refinement of toxins and delivery systems were responsible for its enduring psychological impact. During the period before the issue of effective respirators, Charles Cruttwell, an infantry officer, believed that gas undermined a basic survival mechanism. A serviceman subjected to artillery bombardment had few, if any, defensive options, and trusted to luck. However, when he was exposed to cloud gas, Cruttwell argued, it was impossible to evoke the protection of chance - 'if the very air which he breathes is poison, his chance is gone: he is merely a destined victim for the slaughter'. By contrast, shrapnel was tangible. It could be

removed from a wounded soldier's body by a surgical procedure, but no physician could decontaminate a man's lungs, and it was popularly believed that, once toxins had been metabolized, the respiratory system remained damaged for ever.

While the impact of cloud gas could be assessed because of its very novelty, the subsequent development of chemical weapons in the form of shells and mortar bombs made it more difficult to disaggregate their impact on morale from the wider effects of artillery bombardment. Nevertheless, the Allied armies invested heavily in the production of chemical weapons, and had the war continued into 1919 output was planned to increase significantly,¹⁴⁷ which suggested that commanders had identified a particular casualty producing power in gas. While seasoned infantry battalions habituated to gas, it appears to have unnerved units poorly prepared for the rigours of trench warfare. What was not discovered was whether gas could undermine morale at a faster rate than an artillery barrage.

Because gas shells were mixed with high-explosive ordnance, it was difficult to compare the psychological impact of these various weapons on front-line troops. In addition to the deliberate exploitation of surprise and uncertainty, fears evoked by gas owed much to broad cultural themes. Some toxic chemicals, like phosgene, could not be readily detected through the senses and triggered powerful vestigial fears of mysterious threatening forces. They touched on a deep human concern about the risk of being invaded by a potent and unseen force. Chemical weapons were unfamiliar, which created opportunity for rumour and exaggeration. Beliefs about gas often inspired strong emotions that could disrupt the rational evaluation of evidence and the formation of coping mechanisms. Fears may have been intensified because gas was a product of science and cutting-edge technology. Man-made disasters have generally been experienced as more troubling than natural ones. The novelty and scale of chemical weapons during the First World War were such that they had an enduring impact beyond the veteran population and respiratory medicine. Indeed, in 1928 the US Army physician, Colonel H.L. Gilchrist, wrote, 'the blame for every conceivable sort of ailment has been placed on gas; in fact, there is scarcely a functioning organ of the body whose disturbed action either during or since its participation in [the war] but has had the blame for its erratic performance laid to the door of poison gas'.